arXiv:1704.05780v1 [math.CV] 19 Apr 2017

A REMARK ON PROJECTIVE LIMITS OF FUNCTION SPACES

A. KAMPOUKOU AND V. NESTORIDIS

Abstract. Let $\Omega \subset \mathbb{C}^d$ be an open set and $K_m, m = 1, 2, \dots$ an exhaustion of Ω by compact subsets of Ω. We set $\Omega_m = K_m^o$ and let $X_m(\Omega_m)$ be a topological space of holomorphic functions on Ω_m between $A^\infty(\Omega_m)$ and $H(\Omega_m)$. Then we show that the projective limit of the family $X_m(\Omega_m)$, $m = 1, 2, \dots$, under the restriction maps is homeomorphic and linearly isomorphic to the Fréchet space $H(\Omega)$, idependently of the choice of the spaces $X_m(\Omega_m)$.

1. Introduction

Let Y be a space of holomorhic functions on a domain Ω in $\mathbb{C}^d, d = 1, 2, \dots$. For most of the spaces Y the following holds. If for every bump V of Ω there exists a function f_V in Y which is not holomorphically extendable to V, then there exists a function f in Y not extendable to any V. Moreover, the set of such functions f in Y is dense and G_δ in Y. (see [J-P], [N])

Examples of such spaces are the space $H(\Omega)$ of all holomorphic functions in Ω endowed with the topology of uniform convergence on compacta, the spaces $A^p(\Omega)$ and $H_p^\infty(\Omega)$, Bergman spaces, Hardy spaces etc, provided that Ω satisfies some good assumptions.

Most of these spaces endowed with their natural topology, contain $A^\infty(\Omega)$ and they are contained in $H(\Omega)$: $A^\infty(\Omega) \subset Y \subset H(\Omega)$. Furthermore, the two inclusion maps $i_1 \colon A^\infty(\Omega) \to Y$, $i_2 \colon Y \to H(\Omega)$ with $i_1(f) = f$ and $i_2(w) = w$ are continuous. We denote by $M(\Omega)$ the set of such topological function spaces on Ω.

Let K_m, $m = 1, 2, \dots$, be an exhaustion by compact subsets of the open set $\Omega \subset \mathbb{C}^d$ and $\Omega_m = K_m^o$. Let $X_m(\Omega_m)$ be any topological function space belonging to $M(\Omega_m)$. Let X denote the projective (inverse) limit of the family $X_m(\Omega_m), m = 1, 2, \dots$ where $T_{m,n} : X_m(\Omega_m) \to X_n(\Omega_n), n \leq m$ is the restriction map $f \to f|_{\Omega_n}$.

Then we show that X is a topological vector space homeomorphic and linearly isomorphic to the Fréchet space $H(\Omega)$ of all holomorphic functions

2010 *Mathematics Subject Classification.* 46A13 , 32C18 , 30H.

Key words and phrases. projective limits, holomorphic function, restriction map, Fréchet spaces of holomorphic functions.

on Ω endowed with the topology of uniform convergence on all compact subsets of Ω, independently of the choice of the topological spaces $X_m(\Omega_m)$ from $M(\Omega_m)$. This holds even if the spaces $X_m(\Omega_m)$ are neither complete, nor vector spaces.

2. About projective limits

We need the following particular case of projective limits (see [E]).

Definition 2.1. Let $(Y_n, \mathcal{T}_n), n = 1, 2, \dots$ be a sequence of topological spaces. For $n < m$ let $T_{m,n} \colon Y_m \to Y_n$ be a continuous map and let $T_{n,n}$ be the identity map from Y_n to itself. We assume that for $m \leq n \leq s$ it holds $T_{s,m} = T_{n,m} \circ T_{s,n}$.

Then the projective (or inverse) limit of this family is defined to be the following subset Y of the cartesian product $\prod_{n=1}^{\infty} Y_n$ where

$$Y = \{g = (g_1, g_2, \dots) : \text{ for every } n \leq m \text{ it holds } T_{m,n}(g_m) = g_n\}$$

The topology $\mathcal{T}_Y$ on Y we consider is the relative topology from $\prod_{n=1}^{\infty} Y_n$ endowed with the product topology.

Proposition 2.2. [E] *Under the above notation and assumption a net $g^i = (g_1^i, g_2^i \dots) \in Y, \in I$ converges to $g = (g_1, g_2, \dots) \in Y$ in the topology $\mathcal{T}_Y$ of Y if and only if $g_k^i \to g_k$ in the topology T_k of Y_k for all $k = 1, 2, \dots$.*

3. Some function spaces

Let Ω be an open subset of $\mathbb{C}^d$, $d \geq 1$. We denote by $H(\Omega)$ the space of holomorphic functions on Ω endowed with the topology of the uniform convergence on compacta. This is a Fréchet space with seminorms $\|f\|_m = \sup_{z \in K_m} |f(z)|, m = 1, 2, \dots$ where K_m is an exhaustion of Ω by compact subsets of Ω; for instance we can take (see [R], [G])

$$K_m = \left\{ z \in \Omega : \|z\| \leq m \text{ and } dist(z, \Omega^c) \geq \frac{1}{m} \right\}$$

In what follows we consider that G is a bounded open subset of $\mathbb{C}^d$, although the spaces we will consider, can also be defined for an unbounded open set G, with some modifications in the definition of their natural topology.

The space $A^{\infty}(G)$ is defined as the set of holomorphic functions on G such that each mixed partial derivative of f extends continuously on the closure $\overline{G}$ of G. The natural topology of $A^{\infty}(G)$ is defined by the seminorms

$sup_{z \in G} |\dfrac{\partial^{a_1 + \dots + a_d} f}{\partial {z_1}^{a_1} \dots \partial {z_d}^{a_d}}(z)|, a_1, \dots, a_d \in \{0, 1, 2, \dots\}$, where $z = (z_1, \dots, z_d)$.
This is also a Fréchet space .

Next we are interested in spaces $X(G)$ containing $A^\infty(G)$ and contained in $H(G)$, i.e. $A^\infty(G) \subset Y \subset H(G)$. We also require that the injections are continuous; that is, if a net $f^i \in A^\infty(G)$, $i \in I$, converges in the topology of $A^\infty(G)$ to a function $f \in A^\infty(G)$, then $f^i \to f$ in the topology of $X(G)$, as well; and if a net $w^i \in X(G)$, $i \in I$, converges to $w \in X(G)$ in the topology of $X(G)$ then $w^i \to w$ in the topology of $H(G)$. We denote by $M(G)$ the set of all topological spaces as above.

Such spaces are the spaces $A^p(G)$ consisted of $f \in H(G)$ of which every mixed partial derivative of order less than or equal to p extends continuously on $\overline{G}$ and the spaces $H^p_\infty(G)$ consisted of $f \in H(G)$ of which every mixed partial derivative of order less than or equal to p is bounded on $\overline{G}$. The above two spaces are Banach spaces, provided $p < +\infty$ and Fréchet spaces if $p = +\infty$. Their topologies are defined by the seminorms

$$sup_{z\in G}|\frac{\partial^{a_1+\cdots+a_d} f}{\partial {z_1}^{a_1} \dots \partial {z_d}^{a_d}}(z)|, a_1 + \cdots + a_d \leq p, \text{ where } z = (z_1, \dots, z_d).$$

Another example of such a space is the p-Bergman space, $0 < p < +\infty$, containing all holomorphic functions f on G, such that $\int_G |f|^p d\nu < \infty$, where $d\nu$ denotes the Lebesgue measure on $G \subset \mathbb{C}^d$. (see[D-S])

Other examples are Hardy spaces on the disc or the ball or on more general domains (see [D] , [R2] [S]) , as well as subclasses of the Nevanlinna class , BMOA, VMOA, Dirichlet spaces on the disc etc.

All the previous spaces are complete spaces . However, we can also consider non complete spaces containing $A^\infty(G)$ and contained in $H(G)$ such that the two injections are continuous. For instance, consider the set $Y = A^\infty(G)$ with the relative topology induced by $H(G)$; that is, for $f_n, f \in A^\infty(G)$ the convergence in the new space is the uniform convergence on compacta $f_n \to f$. This space contains $A^\infty(G)$, is contained in $H(G)$ and the injections $i_1 : A^\infty(G) \to Y$, $i_2 \colon Y \to H(G)$ with $i_1(f) = f$ and $i_2(g) = g$ are continuous where $A^\infty(G)$ and $H(G)$ are endowed with their natural topologies and Y with the relative topology from $H(G)$. But this space is not complete, in general. For instance, if $(\overline{G})^o = G$ or if G is a bounded, open subset of $\mathbb{C}$, then $A^\infty(G)$ with the relative topology induced by $H(G)$ is not complete.

It is also possible that a space belongs to $M(G)$ although it is not even a vector space. To give such an example we consider the set

$$Y = A^\infty(G) \cup \{f \colon G \to \mathbb{C} \text{ holomorphic such that } |f(z)| < 1, \text{ for all } z \in G\},$$

endowed with the topology of uniform convergence on G. Then $A^\infty(G) \subset Y \subset H(G)$ and if $f_n, f \in A^\infty(G)$ are such that $f_n \to f$ in the topology

of $A^\infty(G)$, it follows trivially that $f_n \to f$ in the topology of Y. Also if $\phi_n, \phi \in Y$ are such that $\phi_n \to \phi$ in the topology of Y, then obviously the convergence is uniform on each compact subset of G, as well.

4. The result

We start with the following definition

Definition 4.1. Let G be a bounded open subset of Ω^d. We denote by $M(G)$ the set of all topological spaces $X(G)$ satisfying $A^\infty(G) \subset X(G) \subset H(G)$ and such that the two injections $i_1 \colon A^\infty(G) \to X(G)$, $i_1(f) = f$ and $i_2 \colon X(G) \to H(G)$, $i_2(g) = g$ are continuous; that is, if a net $f^i \in A^\infty(G)$, $i \in I$, converges to $f \in A^\infty(G)$, in the topology of $A^\infty(G)$, then $f^i \to f$ in the topology of $X(G)$, as well. Moreover, if a net $w^i \in X(G)$ converges to $w \in X(G)$,in the topology of $X(G)$, then $w_i \to w$ uniformly on compacta of G.

Let Ω be an open subset of C^d, bounded or unbounded. Let $K_m, m = 1, 2, \dots$ be an exhausting family of compact subsets of Ω; that is, $K_m \subset K_{m+1}^o$ and $\cup_{m=1}^\infty K_m = \Omega$ (see [R]),([G]). We set $\Omega_m = K_m^o$; then $\Omega_m \subset \overline{\Omega_m} \subset \Omega_{m+1}$. For each $m = 1, 2, \dots$ let $X_m(\Omega_m)$ be a space in $M(\Omega_m)$.

For $n \leq m$ we consider the restriction map $T_{m,n} \colon X_m(\Omega_m) \to X_n(\Omega_n)$ given by $T_{m,n}(f) = f|_{\Omega_n}$, for every $f \in X_m(\Omega_m)$. These maps are well defined and continuous. Indeed, $X_m(\Omega_m) \subset H(\Omega_m)$ and $\overline{\Omega_n} \subset \Omega_m$; thus, for every $f \in X_m(\Omega_m)$, it follows that $T_{m,n}(f) = f|_{\Omega_n} \in A^\infty(\Omega_n) \subset X_n(\Omega_n)$, provided that $n < m$. Also if a net $f^i \in X_m(\Omega_m)$ converges to $f \in X_m(\Omega_m)$, in the topology of $X_m(\Omega_m)$, then by assumption, it converges uniformly on compacta of Ω_m. Weierstrass theorem yields uniform convergence on compacta of every mixed partial derivative. Since $\overline{\Omega_n}$ is a compact subset of Ω_m, it follows that $f^i \,|_{\Omega_n} \to f|_{\Omega_n}$ in the topology of $A^\infty(\Omega_n)$. By assumption, this implies $f^i \to f$ in the topology of $X_n(\Omega_n)$. Therefore, $T_{m,n} : X_m(\Omega_m) \to X_n(\Omega_n)$ is continuous. One can also very easily verify that $T_{m,n} = T_{s,n} \circ T_{m,s}$, for all $n \leq s \leq m$. Therefore, we can define the projective limit X, which is a topological space (see [E]).

Theorem 4.2. *Under the above assumptions and notation the topological space X is homeomorphic to $H(\Omega)$.*

Proof. Let $f \in H(\Omega)$; then $f|_{\Omega_m} \in A^\infty(\Omega_m) \subset X_m(\Omega_m)$. Thus, if we set $g_m = f|_{\Omega_m}$ and $g = (g_1, g_2, \dots) = g(f)$, then $g \in \prod_{m=1}^\infty X_m(\Omega_m)$. For $n \leq m$ one can easily verify that $T_{m,n}(g_m) = g_n$. Therefore, $g \in X$. In this way we have defined a map $\Phi \colon H(\Omega) \to X$, where $\Phi(f) = g(f)$.

This map is one to one. Indeed, if $g(f) = g(w)$ it follows that $g|_{\Omega_m} = w|_{\Omega_m}$ for all $m = 1, 2, \dots$. Since $\cup_m \Omega_m = \Omega$, it follows that $f(z) = w(z)$ for all $z \in \Omega$; thus $f = w$. This proves that the map Φ is one to one.

The map is also onto. Indeed, if $g = (g_1, g_2, \dots) \in X$, then for every $z \in \Omega$ there exists a least natural number $n(z)$ so that $z \in \Omega_m$, for all $m \geq n(z)$, because $\Omega_m \subset \Omega_{m+1}$ and $\cup_m \Omega_m = \Omega$. Futher, since $T_{m,n(z)} g_m = g_{n(z)}$, it follows that $g_m(z) = g_{n(z)}(z)$, for all $m \geq n(z)$. We set $f(z) = g_{n(z)}(z)$. We will show that this function is holomorphic on Ω. Indeed, let $z_0 \in \Omega$; then $z_0 \in \Omega_{n(z_0)}$. Since $\Omega_{n(z_0)}$ is open, there exists a ball $B(z_0, r)$ contained in $\Omega_{n(z_0)}$. For each $z \in B(z_0, r)$, we have $n(z) \leq n(z_0)$ and $f(z) = g_{n(z)}(z) = g_{n(z_0)}(z)$. But $g_{n(z_0)} \in X_{n(z_0)}(\Omega_{n(z_0)}) \subset H(\Omega_{n(z_0)})$. It follows that f is holomorphic in $B(z_0, r)$. Since z_0 is arbitary in Ω, it follows that $f \in H(\Omega)$. It is immediate that $\Phi(f) = g$. Since g is arbitrary in X, it follows that the map Φ is onto.

Further, Φ is continuous. Indeed, if a net $f^i \in H(\Omega)$, $i \in I$ converges to $f \in H(\Omega)$, uniformly on compacta, by Weierstrass theorem the same holds for every mixed partial derivative. Since $\overline{\Omega_m}$ is a compact subset of Ω it follows that $f^i \mid_{\Omega_m} \to f|_{\Omega_m}$ in the topology of $A^\infty(\Omega_m)$. By assumption this implies the convergence $f^i|_{\Omega_m} \to f \mid_{\Omega_m}$ in the topology of $X_m(\Omega_m)$. According to Proposition 2.2, it follows $\Phi(f^i) \to \Phi(f)$ in the topology of X. Thus, the map Φ is continuous.

Finally, we will show that Φ^{-1} is also continuous. Suppose that $\Phi(f^i) = (f^i|_{\Omega_1}, f^i|_{\Omega_2}, \dots) \to \Phi(f) = (f|_{\Omega_1}, f|_{\Omega_2}, \dots)$ in the topology of X. According to Proposition 2.2 this implies $f^i|_{\Omega_m} \to f|_{\Omega_m}$ for all $m = 1, 2, \dots$ in the topology of $X_m(\Omega_m)$, which, by assumption, implies the uniform convergence on compacta of Ω_m. We will show that $f^i \to f$ uniformly on each compact set $K \subset \Omega$. Let K be such a compact set. Since $K \subset \Omega = \cup_{m=1}^{\infty} \Omega_m$ and the open sets Ω_m satisfy $\Omega_m \subset \Omega_{m+1}$, by compactness of K it follows that there exists m_0 such that $K \subset \Omega_{m_0}$. But $f^i|_{\Omega_{m_0}} \to f|_{\Omega_{m_0}}$ uniformly on each compact of Ω_{m_0}; in particular on K. Thus, $f^i \to f$ uniformly on each compact set $K \subset \Omega$ and therefore, $f^i \to f$ in the topology of $H(\Omega)$. Thus, Φ^{-1} is continuous, as well and Φ is a homeomorphism. This completes the proof. □

Remark 4.3. It is well known and easy to verify that the denumerable projective limit of Fréchet spaces is also a Fréchet space. In our case X is homeomorphic to the Fréchet space $H(\Omega)$, although the spaces $X_m(\Omega_m)$ may not be Fréchet spaces. Furthermore, independently of the fact that the spaces $X_m(\Omega_m)$ are topological vector spaces or they do not have linear

structure, one can easily verify that X is always a topological vector space and that the map Φ is a linear isomorphism.

Remark 4.4. Let us consider the compact set

$$K_t = \left\{ z \in \Omega \colon |z| \leq t, dist(z, \Omega^c) \geq \frac{1}{t} \right\}, \, t \geq t_0,$$

for a convenient $t_0 > 0$ such that $K_{t_0}^o \neq \emptyset$. Then for $t < s$ it follows $K_t \subseteq K_s^o$ and $\cup_{t \geq t_0} K_t = \Omega$. We set $\Omega_t = K_t^o$ and let $X_t(\Omega_t)$ be any element of $M(\Omega_t)$. One can prove that the projective limit of the family $X_t(\Omega_t)$ as $t \to +\infty$, where $T_{s,t}$ is the restriction map $f \to f|_{\Omega_t}$, $t < s$, is homeomorphic and linearly isomorphic to $H(\Omega)$. The proof is similar to that of Theorem 4.2.

Acknowledgement. We would like to thank M.Fragoulopoulou for her interest in this work.

References

[D] P.L. Duren, *Theory of H^p spaces*, Academic Press, N.Y. 1970.

[D-S] P. Duren, A. Schuster, *Bergman spaces*, Mathematical Surveys and Monographs vol.100, AMS.

[E] R. Engelking, *General Topology*, PWN, Warszawa, 1977.

[G] P.M. Gauthier, *Lectures on Several Complex Variables*, Birkhäuser, Heidelberg, N.Y., Dordrecht, London.

[J-P] M. Jarnicki , P. Pflug, *Extension of Holomorphic Functions*, De Gruyter expositions in Mathematics 34, Berlin, N.Y. 2000.

[N] V. Nestoridis, *Domains of Holomorphy*, arXiv:1701.00734.

[R] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, co N.Y., 1966.

[R2] W. Rudin, *Function Theory in the Unit Ball of* $\mathbb{C}^n$, Springer-Verlag, Berlin Heidelberg, 2008.

[S] E. Stein, *Boundary Behavior of Holomorphic Functions of Several Complex Variables*, Math. Notes 11, Princeton University Press, 2016.

Department of Mathematics, National and Kapodistrian University of Athens, Panepistemiopolis, 15784 Athens, Greece

E-mail address: kampouk@math.uoa.gr

E-mail address: vnestor@math.uoa.gr